







pronadisa
Micro & Molecular Biology







Chromogenic Culture Media

knowledge. partnership. future.


CULTURE MEDIA FOR GENERAL USE

| | | | | | | | |
|--|---|--|---|--|---|---|---|
|  Cat. No. 1351 |  Cat. No. 1340 |  Cat. No. 1322 |  Cat. No. 1588 |  Cat. No. 493 |  Cat. No. 2034 |  Cat. No. 2076 |  Cat. No. 2096 |
| TBX Chromogenic Agar According to ISO 16649 | E.coli Coliforms Chromogenic Agar | Salmonella Chromogenic Agar If desired, the supplement Salmonella Chromogenic Agar (Cat. 6043) can be added | E.coli 0157:H7 Chromogenic Agar Base Supplements required (Cat. 6064) | Pseudomonas Chromogenic Agar | Vibrio Chromogenic Agar | Staphylococcus Aureus Chromogenic Agar | Lauryl Sulfate Chromogenic Agar |
| APPLICATION Selective medium for the presumptive detection and enumeration of <i>Escherichia coli</i> | APPLICATION Selective medium for the simultaneous detection of <i>E.coli</i> and other coliforms | APPLICATION For the isolation of <i>Salmonella</i> spp | APPLICATION Selective and differential medium for the detection of <i>E.coli</i> 0157:H7 | APPLICATION For the rapid isolation of Pseudomonas species | APPLICATION For the isolation and detection of <i>Vibrio cholerae</i> , <i>Vibrio parahaemolyticus</i> and <i>Vibrio alginolyticus</i> | APPLICATION For the detection and differentiation different species of staphylococci | APPLICATION For the simultaneous detection of total coliforms and <i>E.coli</i> by the fluorogenic procedure |
| INCUBATION 18-24 hrs at 37°C and 44°C | INCUBATION 18-24 hrs at 35 ± 2°C for total coliforms or at 44 ± 2°C for fecal coliforms | INCUBATION 18-24 hrs at 35± 2°C | INCUBATION 18-24 hrs at 35± 2°C | INCUBATION 24-48 hrs at 35± 2°C | INCUBATION 24-48 hrs at 35 ± 2°C | INCUBATION 24-48 hrs at 35± 2°C | INCUBATION 18-24 hrs at 35± 2°C |

CULTURE MEDIA FOR CLINICAL USE

| | | | | | | |
|---|--|--|--|--|--|---|
|  Cat. No. 1382 |  Cat. No. 1424 |  Cat. No. 1423 |  Cat. No. 1498 |  Cat. No. 2062 |  Cat. No. 2077 |  Cat. No. 2063 |
| Candida Chromogenic Agar | Urinary Tract Infections Chromogenic Agar (UTIC) | MRSA Chromogenic Agar Base Supplements required (Cat. 6069) | MRSA Chromogenic Agar Base, Modified Supplements required (Cat. 6069) | ESBL Chromogenic Agar Base Supplements required (Cat. 6042) | Vancomycin Resistant Enterococcus Chromogenic Agar | KPC Chromogenic Agar Base Supplements required (Cat. 6044) |
| APPLICATION Differential and selective medium for the isolation and quick identification of presumptive <i>Candida</i> spp | APPLICATION For the presumptive detection and differentiation of organisms causing urinary tract infections | APPLICATION For the presumptive detection of Methicillin resistant <i>Staphylococcus aureus</i> | APPLICATION For the presumptive detection of Methicillin resistant <i>Staphylococcus aureus</i> and <i>Staphylococcus epidermidis</i> | APPLICATION For the detection of Gram negative bacteria producing extended Spectrum β-lactamase | APPLICATION For the detection of vancomycin-resistant enterococci | APPLICATION For the detection of Gram negative bacteria with a reduced susceptibility to most of the carbapenem agents |
| INCUBATION 24, 48 and 72 hrs at 30-37°C | INCUBATION 18-24 hrs at 35±2°C | INCUBATION 18-24 hrs at 35± 2°C | INCUBATION 24-48 hrs at 35± 2°C | INCUBATION 18-24 hrs at 35± 2°C | INCUBATION 18-24 hrs at 35 ± 2°C | INCUBATION 18-24 hrs at 35± 2°C |

CULTURE MEDIA FOR INDUSTRY

| | | | | | | | |
|---|--|--|---|---|--|---|---|
|  Cat. No. 2018 |  Cat. No. 1465 |  Cat. No. 1345 |  Cat. No. 1412 |  Cat. No. 2050 |  Cat. No. 1446 |  Cat. No. 1585 |  Cat. No. 2080 |
| E.coli-Enterobacteria Chromogenic Agar | Lauryl Sulfate Chromogenic Broth <i>E.coli</i> is detected by UV light. Indol reaction can be detected | Listeria Chromogenic Agar Base According to ISO 11290. Supplements required (Cat. 6040 and 6031) | m-EI Chromogenic Agar Base Nalidixic acid required | m-EI Chromogenic Agar Base, Modified Nalidixic acid required | Enterobacter Sakazakii Chromogenic Isolation Agar (ESIA) According to ISO 22964 | Standard Methods Chromogenic Agar (PCA) | E.Coli Coliforms Chromogenic Agar (CCA) According to ISO 9308-1 |
| APPLICATION Differential medium for the simultaneous detection of <i>E.coli</i> and Enterobacteria | APPLICATION For the simultaneous detection of <i>E.coli</i> and total coliforms by fluorogenic procedure | APPLICATION Selective medium for the detection and enumeration of <i>Listeria monocytogenes</i> | APPLICATION For the detection and enumeration of <i>Enterococcus</i> through the single step membrane filtration technique | APPLICATION Selective medium for the isolation and differentiation of <i>Enterococcus faecalis</i> and <i>Enterococcus faecium</i> | APPLICATION For the isolation of presumptive <i>Enterobacter sakazakii</i> (<i>Cronobacter sakazakii</i>) | APPLICATION For total microbial plate count | APPLICATION Selective medium for the simultaneous detection of <i>E.coli</i> and other coliforms |
| INCUBATION 18-24 hrs at 35±2°C | INCUBATION 18-24 hrs at 35±2°C | INCUBATION 24 hrs at 37°C | INCUBATION 18-24 hrs at 41 ± 0.5°C | INCUBATION 24-48 hrs at 41 ± 0.5°C | INCUBATION 24 hrs at 44± 1°C | INCUBATION 18-48 hrs at 32± 2°C | INCUBATION 21± 3 hrs at 36 ± 2°C |



For more information contact:

Laboratorios Conda  
Phone: +34 91 761 02 00
Mail: export@condalab.com

Product Packaging Sizes:

*all Media available in Bulk Packs

CULTURE MEDIA FOR GENERAL USE

TBX Chromogenic Agar (ISO 16649-2)
E.coli Coliforms Chromogenic Agar
Salmonella Chromogenic Agar
E.coli O157:H7 Chromogenic Agar Base
Pseudomonas Chromogenic Agar
Vibrio Chromogenic Agar
Staphylococcus Aureus Chromogenic Agar
Lauryl Sulfate Chromogenic Agar

Dehydrated Culture Media 500 g/ 100 g
Dehydrated Culture Media 500 g/ 100 g

Cat. N° 1151
Cat. N° 1340
Cat. N° 1122
Cat. N° 1588
Cat. N° 1493
Cat. N° 2054
Cat. N° 2076
Cat. N° 2096

Supplement for Cat. N° 1122

Cat. N° 6043

Supplement for Cat. N° 1588

Cat. N° 6064

CULTURE MEDIA FOR CLINICAL USE

Candida Chromogenic Agar
Urinary Chromogenic Infections Chromogenic Agar (UTIC)
MRSA Chromogenic Agar Base
MRSA Chromogenic Agar Base, Modified
ESBL Chromogenic Agar Base
KPC Chromogenic Agar Base
Chromogenic Agar For Vancomycin-resistant Enterococcus(VRE)

Dehydrated Culture Media 500 g/ 100 g
Dehydrated Culture Media 500 g/ 100 g

Cat. N° 1382
Cat. N° 1424
Cat. N° 1423
Cat. N° 1498
Cat. N° 2062
Cat. N° 2063
Cat. N° 2077

Supplement for Cat. N° 1423

Cat. N° 6069

Supplement for Cat. N° 1498

Cat. N° 6069

Supplement for Cat. N° 2062

Cat. N° 6042

Supplement for Cat. N° 2063

Cat. N° 6044

CULTURE MEDIA FOR INDUSTRY

E.coli-Enterobacteria Chromogenic Agar
Lauryl Sulfate Chromogenic Broth
Listeria Chromogenic Agar Base (ISO 11290)
m-EI Chromogenic Agar Base
m-EI Chromogenic Agar Base, Modified
Enterobacter Sakazakii Chromogenic Isolation Agar (ESIA) (ISO 22964)
Standard Methods Chromogenic Agar (PCA)
E.Coli Coliforms Chromogenic Agar (CCA) (ISO 9308-1)

Dehydrated Culture Media 500 g/ 100 g
Dehydrated Culture Media 500 g/ 100 g

Cat. N° 2018
Cat. N° 1465
Cat. N° 1345
Cat. N° 1412
Cat. N° 2050
Cat. N° 1446
Cat. N° 1585
Cat. N° 2080

Supplements for Cat. N° 1345

Cat. N° 6031/
N° 6040

Kovacs Reagent for Cat. 1465

Cat. N° 5205



ESTABLISHED 1960

pronadisa
Micro & Molecular Biology

C/ Forja, 9
28850 - Torrejón de Ardoz,
Madrid - SPAIN

Phone. +34 91 761 02 00
Fax +34 91 656 82 28